Long induced paths in $K_{s,s}$ -free graphs

Zach Hunter *

Aleksa Milojević*

Benny Sudakov *

István Tomon[†]

Abstract

More than 40 years ago, Galvin, Rival and Sands showed that every $K_{s,s}$ -free graph containing an n-vertex path must contain an induced path of length f(n), where $f(n) \to \infty$ as $n \to \infty$. Recently, it was shown by Duron, Esperet and Raymond that one can take $f(n) = (\log \log n)^{1/5-o(1)}$. In this note, we give a short self-contained proof that a $K_{s,s}$ -free graphs with an n-vertex path contains an induced path of length at least $(\log \log n)^{1-o(1)}$, which comes closer to the best known upper bound $O((\log \log n)^2)$.

1 Introduction

In 1982, Galvin, Rival and Sands [7] showed that every infinite graph G with a Hamilton path has either arbitrarily long induced paths or it contains the infinite half-graph as a subgraph. Motivated by this result, they asked the following finitary problem: given a graph G that does not contain the complete bipartite graph $K_{s,s}$, and which contains a path on n vertices, how long of an induced path must G contain? They gave a proof showing that every such G contains an induced path of length at least $\Omega((\log \log \log n)^{1/3})$ (here and later, the $\Omega(.), O(.)$, and o(.) notations assume that s is a fixed constant).

This statement was rediscovered some 30 years later, with worse quantitative bounds, by Atminas, Lozin and Razgon [1], in the context of parameterized complexity of the biclique problem. Very recently, Duron, Esperet and Raymond [3] improved upon the bounds of [7], showing that a $K_{s,s}$ -free graph G with an *n*-vertex path must contain an induced path of length at least $(\log \log n)^{1/5-o(1)}$. On the other hand, a surprising result of Defrain and Raymond [4] shows the existence of 2-degenerate *n*-vertex graphs with a Hamilton path but no induced paths of length $\Omega((\log \log n)^2)$. Note that 2-degenerate graphs are also $K_{3,3}$ -free. The purpose of our paper is to give a very short proof of the following improved lower bound.

Theorem 1.1. Let $n \ge s \ge 2$ be positive integers and let G be a $K_{s,s}$ -free graph containing a path on n vertices. Then G contains an induced path of length at least $\Omega(\frac{\log \log n}{\log \log \log n})$.

Determining the length of longest induced paths in hereditary classes of graphs has been the subject of many recent papers, see [3] for an excellent overview of the results related to this problem. A result of Nešetřil and Ossona de Mendez [9] addresses the class of d-degenerate graphs, showing that a d-degenerate graph must contain an induced path of length $\Omega(\log \log n/\log d)$. As mentioned above, this bound cannot be improved beyond $(\log \log n)^2$ already for d = 2 by a construction of Defrain and Raymond [4].

More generally, the study of long induced paths in graphs with structural properties fits well into the broader line of research about induced subgraphs of $K_{s,s}$ -free or K_s -free graphs. The assumption that the host graph G is $K_{s,s}$ -free or K_s -free is a natural one, since it forbids the host graph from being a complete graph, which contains no nontrivial induced subgraphs. An example of this type of problem is the old question of Erdős, Saks and Sós [5], which asks to determine the maximum size of an induced tree in a connected K_s -free graph. This problem exhibits an interesting transition between s = 3 and $s \ge 4$, since the largest induced tree in a K_3 -free graph is of size $\Omega(\sqrt{n})$, while for $s \ge 4$ there are K_s -free graphs without induced trees on more than $O(\log n)$ vertices, see [6].

^{*}Department of Mathematics, ETH Zürich, Switzerland. Email: {zach.hunter, aleksa.milojevic, benjamin.sudakov}@math.ethz.ch. Research supported in part by SNSF grant 200021-228014.

[†]Umeå University, *e-mail*: **istvantomon@gmail.com**, Research supported in part by the Swedish Research Council grant VR 2023-03375.

The systematic study of induced subgraphs of $K_{s,s}$ -free graphs was initiated by the authors of this note in the recent paper [8], posing the following general Turán-type question. Given a fixed graph H, what is the maximum number of edges in a $K_{s,s}$ -free graph G on n vertices which does not contain H as an induced subgraph? If s and H are fixed, it is shown for several interesting classes of graphs, such as trees or cycles, that this maximum number of edges cannot exceed the usual extremal number ex(n, H) by more than a constant factor depending on s and H, and it was conjectured that this phenomenon holds for all bipartite graphs H.

Finally, let us mention that the study of $K_{s,s}$ -free graphs without certain induced subgraphs also comes up naturally in geometry. Namely, incidence or intersection graphs of various geometric objects often avoid certain induced subgraphs for geometric reasons, and thus the maximum number of edges needed to find a $K_{s,s}$ can be studied through the methods of graph theory. For more detailed discussions of this subject, see the excellent survey of Smorodinsky [10].

2 Long induced paths

In this section, we prove Theorem 1.1 after a few preliminary lemmas. Given a graph G and $v \in V(G)$, N(v) denotes the neighbourhood of v.

Lemma 2.1. Let $d, s \ge 2$ be integers and $t \ge 1$ be a real number satisfying $t \le d^{1/s}/s$. If G is a $K_{s,s}$ -free graph of minimum degree at least d and v is a vertex of G, then less than st vertices $u \in V(G)$ satisfy $|N(u) \cap N(v)| \ge \frac{4}{t}|N(v)|$.

Proof. Suppose for the sake of contradiction that that there exists a set U of size |U| = st such that for every $u \in U$, we have $|N(u) \cap N(v)| \ge \frac{4}{t}|N(v)|$. Consider the bipartite subgraph H of G induced between the sets U and $N(v)\setminus U$.

Each vertex of U has at least $\frac{4}{t}|N(v)| - |U| \ge \frac{2}{t}|N(v)|$ neighbours in $N(v)\setminus U$. Here, we used that $|U| = st \le \frac{2}{t}d \le \frac{2}{t}|N(v)|$, which follows from the the assumptions $t \le d^{1/s}/s$ and $s \ge 2$. Hence, the number of edges in H is at least $|U| \cdot \frac{2}{t}|N(v)|$. On the other hand, the Kővári-Sós-Turán theorem implies that in a $K_{s,s}$ -free bipartite graph with parts of size m = |U| and $n = |N(v)\setminus U|$, one can have at most $(s-1)^{1/s}mn^{1-1/s} + (s-1)n$ edges (see e.g. Theorem 2.2 in Chapter VI of [2]). In our case,

$$\frac{2}{t}|U||N(v)| \le e(U, N(v)\backslash U) < s|U||N(v)|^{1-1/s} + s|N(v)|.$$
(1)

But $\frac{1}{t}|U||N(v)| \ge s|U||N(v)|/d^{1/s} \ge s|U||N(v)|^{1-1/s}$, since $|N(v)| \ge d$ and $\frac{1}{t}|U||N(v)| = s|N(v)|$. This contradicts (1), finishing the proof.

Lemma 2.2. Let G be a $K_{s,s}$ -free graph of minimum degree at least d. Then G contains an induced path of length at least $d^{1/2s}/2s$.

Proof. Fix $k = \lceil \frac{1}{2s} d^{1/2s} \rceil$ and assume $k \ge 4$ since the statement is trivial otherwise. We pick a random walk v_1, \ldots, v_k according to the stationary distribution. Namely, the starting vertex v_1 is chosen at random from V(G) such that $\mathbb{P}[v_1 = w] = \frac{\deg w}{2e(G)}$ for all $w \in V(G)$, and each subsequent vertex v_{i+1} is chosen uniformly at random among the neighbours of v_i . Note that this choice ensures that (v_i, v_j) has the same distribution as (v_j, v_i) for any $1 \le i < j \le k$ (this follows since the walk (v_1, \ldots, v_k) has the same distribution as its reverse (v_k, \ldots, v_1)). We show that the walk v_1, \ldots, v_k is an induced path with positive probability.

Define an auxiliary directed graph H on the vertex set V(G), where $v \to u$ if $|N(u) \cap N(v)| \ge \frac{4s}{d^{1/s}} |N(v)|$. Lemma 2.1 applied with $t = d^{1/s}/s$ shows that the outdegree of a vertex v in H is at most $d^{1/s}$. Next, we estimate the probability that there is no edge $v_j \to v_i$ for some $1 \le i < j \le k$ in H, and we denote this event by E. By the union bound, we have

$$\mathbb{P}[\overline{E}] = \mathbb{P}[v_j \to v_i \text{ for some } 1 \le i < j \le k] \le \sum_{1 \le i < j \le k} \mathbb{P}[v_j \to v_i].$$

Recall that the pair (v_i, v_j) has the same distribution as (v_j, v_i) , so $\mathbb{P}[v_j \to v_i] = \mathbb{P}[v_i \to v_j]$. Since v_j is chosen uniformly from $N_G(v_{j-1})$, which is a set of size at least d, and the outdegree of v_i in H is at most $d^{1/s}$, we find that $\mathbb{P}[v_i \to v_j] \leq \frac{d^{1/s}}{d} = d^{1/s-1}$. Thus, $\mathbb{P}[\overline{E}] \leq {k \choose 2} d^{1/s-1} \leq k^2 d^{1/s-1}$.

The next step is to estimate the probability that v_1, \ldots, v_k is an induced path. The walk v_1, \ldots, v_k is an induced path if and only if there are no edges between v_i and v_j for i < j - 1. Indeed, since $k \ge 4$, the latter condition implies that the vertices v_1, \ldots, v_k are all distinct. Fixing some $1 \le i < j - 1 \le k - 1$, our main observation is that $\mathbb{P}[v_i v_j \in E(G)|v_{j-1} \nleftrightarrow v_i] \le 4sd^{-1/s}$. Indeed, if $v_{j-1} \nleftrightarrow v_i$ in H, then at most $\frac{4s}{d^{1/s}}|N(v_{j-1})|$ neighbours $v_j \in N(v_{j-1})$ form an edge of G with v_i . As a consequence, we have

$$\mathbb{P}[v_i v_j \in E(G) \text{ and } E] \leq \mathbb{P}[v_i v_j \in E(G)|E] \leq 4sd^{-1/s}$$

We now complete the proof. If v_1, \ldots, v_k is not an induced path in G, then either E does not happen, or there exist some i < j - 1 for which $v_i v_j \in E(G)$ and $v_{j-1} \nleftrightarrow v_i$ in H. Since the number of such pairs (i, j) is bounded by $\binom{k}{2}$, and we have $k = \frac{1}{2s} d^{1/2s}$, $s \ge 2$, we conclude that

$$\mathbb{P}[v_1, \dots, v_k \text{ is not an induced path in } G] \leq \mathbb{P}[\overline{E}] + \sum_{1 \leq i < j-1 \leq k-1} \mathbb{P}[v_i v_j \in E(G) \text{ and } E]$$
$$\leq k^2 d^{1/s-1} + 4s \binom{k}{2} d^{-1/s} \leq \frac{1}{4s^2} d^{2/s-1} + \frac{1}{2s} < 1.$$

Finally, we give the proof of the following lemma of Nešetřil and Ossona de Mendez [9] for completeness. Recall that a graph G is d-degenerate if every subgraph of G contains a vertex of degree $\leq d$.

Lemma 2.3. Let G be a d-degenerate graph on n vertices containing a Hamilton path. Then G contains an induced path of length $\Omega(\frac{\log \log n}{\log d})$.

Proof. We start with a simple claim.

Claim 2.4. If G is an acyclic directed graph of maximum outdegree d on n vertices with a directed Hamilton path, then G contains an induced directed path of length at least $\Omega(\frac{\log n}{\log d})$.

Proof. Let v_1 be the first vertex of the Hamilton path, and run the breadth-first search algorithm (BFS algorithm) starting at v_1 . Since every vertex of G can be reached from v_1 by a directed path, the algorithm explores the whole graph and constructs a spanning tree, where each node has at most d children. This tree has n vertices and so it must have depth at least $\Omega(\frac{\log n}{\log d})$. Moreover, by simple properties of the BFS algorithm and the acyclic property of G, each directed path from the root to a leaf is induced, thus giving us an induced path of the desired length.

Denote the vertices of G by v_1, \ldots, v_n in the order they appear on the Hamilton path. Since G is a d-degenerate graph, its edges can be directed such that the outdegree of each vertex in the resulting directed graph is at most d. Let G' be the subgraph consisting of the edges of the Hamilton path together with all edges $v_i \to v_j$ with i < j, and direct all the edges of the Hamilton path from v_i to v_{i+1} . Then G'satisfies the conditions of Claim 2.4 (with d + 1 instead of d), so it contains an induced path P of length $|P| \ge \Omega(\frac{\log n}{\log d})$.

Let the vertices of P be v_{i_1}, \ldots, v_{i_k} with $i_1 < \cdots < i_k$. Consider the subgraph of G induced on the vertices of P. Since P is induced in G', if $v_{i_a}v_{i_b} \in E(G)$ for some a < b - 1, then $v_{i_b} \to v_{i_a}$ in G. Reverse the direction of the edges $v_{i_a}v_{i_{a+1}}$ for $a = 1, \ldots, k - 1$, and let the resulting directed graph be P'. Then P' satisfies the conditions of Claim 2.4, so we find a path $P'' \subseteq P'$ which is induced in P', and thus also in G, of length $\Omega(\frac{\log |P|}{\log d}) \ge \Omega(\frac{\log \log n}{\log d})$. This completes the proof.

Now, we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Let $d = (\log \log n)^{2s}$. If G is a d-degenerate graph, then by Lemma 2.3, G contains an induced path of length $\Omega(\frac{\log \log n}{\log d}) = \Omega(\frac{\log \log n}{2s \log \log \log n}) = \Omega(\frac{\log \log n}{\log \log \log n})$, as claimed. If G is not d-degenerate, it contains an induced subgraph G' with minimum degree at least d. Since G' is also $K_{s,s}$ -free, Lemma 2.2 implies that G' contains an induced path of length $d^{1/2s}/2s = \Omega(\log \log n)$.

References

- A. Atminas, V. V. Lozin and I. Razgon, Linear time algorithm for computing a small biclique in graphs without long induced paths, in Algorithm theory—SWAT 2012, 142–152, Lecture Notes in Comput. Sci., 7357, Springer, Heidelberg.
- [2] B. Bollobás, Extremal Graph Theory, Dover Publications, 2004.
- [3] J. Duron, L. Esperet, J.-F. Raymond, Long induced paths in sparse graphs and graphs with forbidden patterns, preprint, arXiv:2411.08685.
- [4] O. Defrain and J.-F. Raymond, Sparse graphs without long induced paths, J. Combin. Theory Ser. B 166 (2024), 30–49.
- [5] P. Erdős, M. E. Saks and V. T. Sós, Maximum induced trees in graphs, J. Combin. Theory Ser. B 41 (1986), 61–79.
- [6] J. Fox, P.-S. Loh, B. Sudakov, Large induced trees in K_r -free graphs, J. Combin. Theory Ser. B 99 (2009), 494–501.
- [7] F. Galvin, I. Rival and B. Sands, A Ramsey-type theorem for traceable graphs, J. Combin. Theory Ser. B 33 (1982), 7–16.
- [8] Z. Hunter, A. Milojević, B. Sudakov and I. Tomon, Kővári-Sós-Turán theorem for hereditary families, preprint, arXiv:2401.10853.
- [9] J. Nešetřil and P. Ossona de Mendez, Sparsity, Springer, 2012.
- [10] S. Smorodinsky, A survey of Zarankiewicz problem in geometry, preprint, arXiv:2410.03702.